

Human IL-3

research grade

10 µg	130-093-908
25 µg	130-093-909
4×25 µg	130-094-193
1000 µg	130-093-911

Contents

1. Description
 - 1.1 Background information
 - 1.2 Applications
2. References

1. Description

Components	Human IL-3, research grade: Purified recombinant human interleukin 3.
Sizes	10 µg, 25 µg, 4×25 µg, 1000 µg.
Biological activity	The ED ₅₀ is ≤1.0 ng/mL* corresponding to a specific activity of ≥1×10 ⁶ I.U./mg.
Primary structure	Single, non-glycosylated polypeptide chain (133 amino acid residues).
Molecular mass	15.1 kDa.
Source	Produced in <i>E. coli</i> .
Product format	Lyophilized from a 0.2 µm filtered sodium chloride solution.
Stabilizer	Mannitol and/or trehalose (HSA, 1000 µg size).
Purity	>97% as determined by SDS-PAGE analysis.
Endotoxin level	Low endotoxin (<1.0 EU/µg cytokine) as determined by Limulus Amebocyte Lysate (LAL) assay.
Storage	Lyophilized Human IL-3, research grade should be stored at -20 °C. The expiration date is indicated on the vial label. Upon reconstitution aliquots should be stored at -20 °C. Avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended to reconstitute lyophilized Human IL-3 with deionized sterile-filtered water to a final concentration of not less than 100 µg/mL.

▲ **Note:** Addition of carrier protein, such as 0.1% bovine serum albumin (BSA) or human serum albumin (HSA) may have stabilizing effects. Further dilutions should be prepared with 1% BSA or HSA in phosphate-buffered saline (PBS).

* The ED₅₀ is determined by proliferation assay using TF-1 cells according to Kitamura *et al.*¹ The proliferation assay was calibrated with the international standard for human IL-3 (NIBSC code 91/510) provided by the WHO/National Institute for Biological Standards and Control.

1.1 Background information

Interleukin 3 (IL-3) is a hematopoietic growth factor which is produced mainly by activated T cells, but is also secreted by other cell types, including mast cells, eosinophils, and keratinocytes. The broad spectrum of biologic activities of IL-3 includes the stimulation of the proliferation and differentiation of immature pluripotent hematopoietic stem cells and various lineage-committed progenitor

cells, leading to the production of most of the major blood cell types. In addition, IL-3 also affects the functional activity of mature mast cells, basophils, eosinophils, and macrophages.

1.2 Applications

Human IL-3 can be used for a variety of applications, including:

- Induction of colony formation from hematopoietic progenitor cells in semi-solid medium *in vitro*, for example, CD34⁺ cells from umbilical cord blood².
- *In vitro* differentiation studies, for example, of B lymphoid progenitors³.
- Cultivation of plasmacytoid dendritic cells.⁴
- Investigation of mast cell or basophil function, for example, basophil interaction with blood vessels⁵.
- Investigation of IL-3-mediated signaling pathways.

Optimal concentration for a specific application should be determined by a dose-response experiment.

2. References

1. Kitamura, T. *et al.* (1989) Establishment and characterization of a unique human cell line that proliferates dependently on GM-CSF, IL-3, or erythropoietin. *J. Cell Physiol.* 140: 323–334.
2. Rossmann, T. *et al.* (2001) Interleukin 3 improves the ex vivo expansion of primitive human cord blood progenitor cells and maintains the engraftment potential of SCID repopulating cells. *Stem Cells* 19: 313–320.
3. Crooks, G. M. *et al.* (2000) IL-3 increases production of B lymphoid progenitors from human CD34⁺CD38⁻ cells. *J. Immunol.* 165: 2382–2389.
4. Tas, S. W. *et al.* (2007) Noncanonical NF-kappaB signaling in dendritic cells is required for indoleamine 2,3-dioxygenase (IDO) induction and immune regulation. *Blood* 110: 1540–1549.
5. Lim, L. H. *et al.* (2006) Stimulation of hman endothelium with IL-3 induces selective basophil accumulation in vitro. *J. Immunol.* 176: 5346–5353.

All protocols and data sheets are available at www.miltenyibiotec.com.

Warranty

The products sold hereunder are warranted only to be free from defects in workmanship and material at the time of delivery to the customer. Miltenyi Biotec GmbH makes no warranty or representation, either expressed or implied, with respect to the fitness of a product for a particular purpose. There are no warranties, expressed or implied, which extend beyond the technical specifications of the products. Miltenyi Biotec GmbH's liability is limited to either replacement of the products or refund of the purchase price. Miltenyi Biotec GmbH is not liable for any property damage, personal injury or economic loss caused by the product.

MACS is a registered trademark of Miltenyi Biotec GmbH.

Unless otherwise specifically indicated, Miltenyi Biotec products and services are for research use only and not for diagnostic or therapeutic use.

Copyright © 2009 Miltenyi Biotec GmbH. All rights reserved.