

Contents

1. Description
 - 1.1 Background information
 - 1.2 Applications
 - 1.3 Recommended antibody dilution
 - 1.4 Reagent requirements
2. General protocol for immunofluorescent staining
3. Examples of immunofluorescent staining with Anti-IgG1 antibodies

1. Description

Components	1 mL monoclonal Anti-IgG1 antibodies, mouse conjugated to various dyes.	
	FITC	130-095-897
	PE	130-095-900
	APC	130-095-902
	Biotin	130-095-879
Clone	X-56 (isotype: rat IgG1).	
Capacity	100 tests or up to 10 ⁹ total cells.	
Product format	Antibodies are supplied in buffer containing stabilizer and 0.05% sodium azide.	
Storage	Store protected from light at 2–8 °C. Do not freeze. The expiration date is indicated on the vial label.	

1.1 Background information

The Anti-IgG1 antibody, mouse is suited for the indirect immunofluorescent staining and flow cytometric analysis of cells stained with primary mouse antibodies of IgG1 isotype. The antibody can further be used for flow cytometric analysis of mouse B cells expressing surface IgG1, for example, memory B cells.

1.2 Applications

- Identification and enumeration of IgG1⁺ cells by flow cytometry or fluorescence microscopy.
- Detection of cells labeled with a primary mouse antibody of IgG1 isotype.

1.3 Recommended antibody dilution

The recommended antibody dilution for all Anti-IgG1 conjugates is **1:11 for up to 10⁷ cells/100 µL** of buffer for labeling of cells and analysis by flow cytometry.

The antibody is suited for staining of formaldehyd-fixed cells.

1.4 Reagent requirements

- Buffer: Prepare a solution containing phosphate-buffered saline (PBS), pH 7.2, 0.5% bovine serum albumin (BSA), and 2 mM EDTA by diluting MACS® BSA Stock Solution (# 130-091-376) 1:20 with autoMACS® Rinsing Solution (# 130-091-222). Keep buffer cold (2–8 °C).
 - ▲ **Note:** EDTA can be replaced by other supplements such as anticoagulant citrate dextrose formula-A (ACD-A) or citrate phosphate dextrose (CPD). BSA can be replaced by other proteins such as mouse serum albumin, mouse serum, or fetal bovine serum (FBS). Buffers or media containing Ca²⁺ or Mg²⁺ are not recommended for use.
- (Optional) FcR Blocking Reagent, mouse (# 130-092-575) to avoid Fc receptor-mediated antibody labeling.
- (Optional) Anti-Biotin antibodies conjugated to, e.g., Anti-Biotin-PE (# 130-090-756) as secondary antibody reagent in combination with Anti-IgG1-Biotin.
- (Optional) Primary mouse antibodies of IgG1 isotype or antibodies for staining of B cells. For more information about antibodies refer to www.miltenyibiotec.com/antibodies.
- (Optional) Propidium Iodide Solution (# 130-093-233) or 7-AAD for flow cytometric exclusion of dead cells without fixation.
- (Optional) Fixation and Dead Cell Discrimination Kit (# 130-091-163) for cell fixation and flow cytometric exclusion of dead cells.

2. General protocol for immunofluorescent staining

▲ Volumes given below are for **up to 10⁷** nucleated cells. When working with fewer than 10⁷ cells, use the same volumes as indicated. When working with higher cell numbers, scale up all reagent volumes and total volumes accordingly (e.g. for 2×10⁷ nucleated cells, use twice the volume of all indicated reagent volumes and total volumes).





▲ For detection of cells labeled with a primary antibody of mouse IgG1 isotype stain cells according to manufacturer's recommendation. After washing proceed with step 2.3.

1. Determine cell number.
2. Centrifuge cell suspension at 300×g for 10 minutes. Aspirate supernatant completely.
3. Resuspend up to 10⁷ nucleated cells per 100 µL of buffer.
4. Add 10 µL of the Anti-IgG1 antibody.
5. Mix well and incubate for 10 minutes in the dark in the refrigerator (2–8 °C).
 - ▲ **Note:** Higher temperatures and/or longer incubation times may lead to non-specific cell labeling. Working on ice requires increased incubation times.
6. Wash cells by adding 1–2 mL of buffer and centrifuge at 300×g for 10 minutes. Aspirate supernatant completely.

7. (Optional) If Anti-IgG1-Biotin was used, resuspend the cell pellet in 100 μ L of buffer, add 10 μ L of anti-biotin antibody, and continue as described in steps 5 and 6.
8. Resuspend cell pellet in a suitable amount of buffer for analysis by flow cytometry or fluorescence microscopy.

3. Examples of immunofluorescent staining with Anti-IgG1 antibodies

Human peripheral blood mononuclear cells (PBMCs) were first labeled with anti-human CD4 (isotype IgG1) and then stained with Anti-IgG1 antibodies conjugated to FITC (A), PE (B), APC (C), or Biotin (D) and analyzed by flow cytometry using the MACSQuant[®] Analyzer. Cell debris and dead cells were excluded from the analysis based on scatter signals and propidium iodide fluorescence.



All protocols and data sheets are available at www.miltenyibiotec.com.

Warnings

Reagents contain sodium azide. Under acidic conditions sodium azide yields hydrazoic acid, which is extremely toxic. Azide compounds should be diluted with running water before discarding. These precautions are recommended to avoid deposits in plumbing where explosive conditions may develop.

Warranty

The products sold hereunder are warranted only to be free from defects in workmanship and material at the time of delivery to the customer. Miltenyi Biotec GmbH makes no warranty or representation, either expressed or implied, with respect to the fitness of a product for a particular purpose. There are no warranties, expressed or implied, which extend beyond the technical specifications of the products. Miltenyi Biotec GmbH's liability is limited to either replacement of the products or refund of the purchase price. Miltenyi Biotec GmbH is not liable for any property damage, personal injury or economic loss caused by the product.

autoMACS, MACS, and MACSQuant are registered trademarks of Miltenyi Biotec GmbH.

Copyright © 2010 Miltenyi Biotec GmbH. All rights reserved.