



Reference list

Brain Tumor Dissociation Kits

Töpfer, K. *et al.* (2015) DAP12-Based Activating Chimeric Antigen Receptor for NK Cell Tumor Immunotherapy. *J. Immunol.*, 194: 3201-3212.

Kusne, Y. *et al.* (2014) Targeting aPKC disables oncogenic signaling by both the EGFR and the proinflammatory cytokine TNF α in glioblastoma. *Sci. Signal* 7: ra75.

Santagata, S. *et al.* (2014) Intraoperative mass spectrometry mapping of an onco-metabolite to guide brain tumor surgery. *Proc. Natl. Acad. Sci. U.S.A.* 111: 11121-11126.

Zeiner, P. S. *et al.* (2014) MIF receptor CD74 is restricted to microglia/macrophages, associated with a M1-polarized immune milieu and prolonged patient survival in gliomas. *Brain Pathol.* 25 (4): 491-504.

Golebiewska, A. *et al.* (2013) Side population in human glioblastoma is non-tumorigenic and characterizes brain endothelial cells. *Brain* 136 (Pt 5): 1462-1475.

Ji, M. *et al.* (2013) Rapid, label-free detection of brain tumors with stimulated raman scattering microscopy. *Sci. Transl. Med.* 5: 201ra119.

Piao, Y. *et al.* (2013) Acquired resistance to anti-VEGF therapy in glioblastoma is associated with a mesenchymal transition. *Clin. Cancer Res.* 19 (16): 4392-4403.

Vinnakota, K. *et al.* (2013) Toll-like receptor 2 mediates microglia/brain macrophage MT1-MMP expression and glioma expansion. *Neuro Oncol.* 15: 1457-1468.

Janiszewska, M. *et al.* (2012) Imp2 controls oxidative phosphorylation and is crucial for preserving glioblastoma cancer stem cells. *Genes Dev.* 26 (17): 1926-1944.


Piao, Y. *et al.* (2012) Glioblastoma resistance to anti-VEGF therapy is associated with myeloid cell infiltration, stem cell accumulation, and a mesenchymal phenotype. *Neuro Oncol.* 14 (11): 1379-1392.

Kijima, N. *et al.* (2011) CD166/activated leukocyte cell adhesion molecule is expressed on glioblastoma progenitor cells and involved in the regulation of tumor cell invasion. *Neuro Oncol.* 14 (10): 1254-1264.

Skog, J. *et al.* (2008) Glioblastoma microvesicles transport RNA and proteins that promote tumour growth and provide diagnostic biomarkers. *Nat. Cell. Biol.* 10 (12): 1470-1476.

Watch this video protocol demonstrating how to isolate microglial cells from adult mouse brain using Miltenyi Biotec's standardized workflow.

The procedure for dissociation, myelin removal, and cell isolation of brain tumors is similar to that for adult mouse brain.



Product	Order no.
Brain Tumor Dissociation Kit (P)	130-095-942
Brain Tumor Dissociation Kit (T)	130-095-939
Myelin Removal Beads II, human, mouse, rat – small size (200 μ L)	130-096-731
Myelin Removal Beads II, human, mouse, rat (4 mL)	130-096-733
Myelin Removal Beads II, human, mouse, rat (2x4 mL)	130-096-433
gentleMACS™ Dissociator	130-093-235
gentleMACS™ Octo Dissociator	130-095-937
gentleMACS™ Octo Dissociator with Heaters	130-096-427

Miltenyi Biotec GmbH | Phone +49 2204 8306-0 | Fax +49 2204 85197 | macs@miltenyibiotec.de | www.miltenyibiotec.com

Miltenyi Biotec provides products and services worldwide. Visit www.miltenyibiotec.com/local to find your nearest Miltenyi Biotec contact.

Unless otherwise specifically indicated, Miltenyi Biotec products and services are for research use only and not for therapeutic or diagnostic use. MACS and the MACS logo are registered trademarks or trademarks of Miltenyi Biotec GmbH.