

Allogeneic transplantation and expansion of CD133 positive cells**Successful transplantation of haploidentically mismatched peripheral blood stem cells using CD133+-purified stem cells.**

Bitan M, Shapira MY, Resnick IB, Zilberman I, Miron S, Samuel S, Ackerstein A, Elad S, Israel S, Amar A, Fibach E, Or R, Slavin S.
Exp Hematol. 2005; 33(6):713-8.

Rapid reconstitution of dendritic cells after allogeneic transplantation of CD133+ selected hematopoietic stem cells.

Bornhauser M, Eger L, Oelschlaegel U, Auffermann-Gretzinger S, Kiani A, Schetelig J, Illmer T, Schaich M, Corbeil D, Thiede C, Ehninger G.
Leukemia. 2005 Jan;19(1):161-5.

Transplantation of a combination of CD133+ and CD34+ selected progenitor cells from alternative donors.

Lang P, Bader P, Schumm M, Feuchtinger T, Einsele H, Fuhrer M, Weinstock C, Handgretinger R, Kuci S, Martin D, Niethammer D, Greil J.
Br J Hematol. 2004 Jan; 124(1):72-79.

Correction of persistent thrombocytopenia by a boost of CD133+ selected stem cells in a patient transplanted for Wiskott-Aldrich syndrome 10 years ago.

Lang P, Klingebiel T, Schumm M, Handgretinger R, Greil J, Beck JF, Niethammer D, Bader P.
Bone Marrow Transplant. 2004 Apr; 33(8):879-80.

Large-scale isolation of CD133+ progenitor cells from G-CSF mobilized peripheral blood stem cells.

Gordon PR, Leimig T, Babarin-Dorner A, Houston J, Holladay M, Mueller I, Geiger T, Handgretinger R.
Bone Marrow Transpl. 2003 Jan; 31(3):17-22.

CD133+ selection is an alternative to CD34+ cell selection for ex-vivo expansion of HSCs.

Kobari L, Giarratana MC, Pflumio F, Izac B, Coulombel L, Douay L.
J Hematother Stem Cell Res. 2001 Apr; 10(2): 273-281.

Abstracts**Boosts with positive selected stem cells can improve haematopoietic recovery after allogeneic transplantation.**

Lang P, Greil J, Schumm M, Ebinger M, Bader P, Schlegel PG, Pfeiffer M, Feuchtinger T, Handgretinger R.
Bone Marrow Transplant. 2006; 37(S1):S67.

Lack of severe acute Graft-versus-Host disease following transplantation of purified CD34+CD133+ cells with T-cell add-back from unrelated donors.

Greil J, Lang P, Bader P, Schlegel PG, Schumm M, Feuchtinger T, Niethammer D.
Blood 2003; 102(11): 490a.

Improved platelet recovery after transplantation of highly purified allogeneic CD133+ stem cells in children.

Lang P, Bader P, Niethammer D, Schumm M, Schwaemmle M, Jaki C, Weinstock C, Kuci S, Feuchtinger T, Handgretinger R, Greil J.
Bone Marrow Transpl. 2003; 31(Suppl1):S795.

Positive enrichment of CD133+(AC133+) cells with the CliniMACS device: comparison to CD34.

Schumm M, Lang P, Kuci S, Greil J, Niethammer D.
Bone Marrow Transpl 2003; 31(Suppl1):S211.

Transplantation of CD133+ selected haploidentical hematopoietic stem cells.

Greil J, Lang PJ, Schumm M, Bader P, Kuci S, Gordon P, Handgretinger R, Niethammer R.
Blood 2002; 100(11): 637a.

Transplantation of human CD133 selected stem cells results in enhanced engraftment and Tcell reconstitution in NOD/SCID mice.

Gordon P, Leimig T, Chen X, Babarin-Dorner A, Holladay M, Houston J, Barfield R, Srivastava K, Handgretinger R
Blood 2002; 100(11): 713a.

Third party donor CD133 selected stem cells in cord blood transplantation**Abstracts****Cord blood transplantation supported by co-infusion of CD133-positive haematopoietic stem cells from a third party donor: preliminary results.**

Krsnik I, Regidor C, Cabrera JR, Forés R, Ruiz E, Bautista G, García-Marco JA, Ojeda E, Sanjuan I, Gil S, Magro E, Vallejo C, Fernandez MN.
Bone Marrow Transplant. 2006; 37(S1): S55.

Selection of CD133+ haematopoietic progenitor cells for use in allogeneic transplantation.

Krsnik I, Regidor C, Forés R, Bautista G, Cabrera JR, Ruiz E, Ojeda E, Gil S, Sanjuan I, García-Marco JA, Vallejo C, Fernandez MN.
Bone Marrow Transplant. 2006; 37(S1): S64.

Autologous transplantation of CD133 positive cells and purging**Immunologic purging of autologous peripheral blood stem cell products based on CD34 and CD133 expression can be effectively and safely applied in half of the acute myeloid leukemia patients.**

Feller N, van der Pol MA, Waaijman T, Weijers GW, Westra G, Ossenkoppele GJ, Schuurhuis GJ.
Clin Cancer Res. 2005 Jul 1; 11(13):4793-801.

Autologous transplantation of CD133 selected hematopoietic progenitor cells in a pediatric patient with relapsed leukemia.

Koehl U, Zimmermann S, Esser R, Sorensen J, Gruttner HP, Duchscherer M, Seifried E, Klingebiel T, Schwabe D.
Bone Marrow Transpl. 2002 Jun;29(11):927-930.

AC133 for tumor cell depletion

Köhl-U
MACS&MORE 2000 ; 4(2):12-13

Abstracts**Selection and autologous transplantation of rhG-CSF mobilized CD133+ cells from peripheral blood in patients with indolent Non Hodgkin's Lymphoma under high-dose chemotherapy.**

Falzetti F, Ballanti S, De Ioanni M, Zugarini V, Piattoni S, Battelli F, Zei T, Iacucci Ostini R, Martelli MF, Tabilio A.
Blood 2002; 100(11): 831a.

Purging of acute myeloid leukemia transplants by positive selection using CD133 or CD34 expression

Schuurhuis GJ, Waaijman T, van der Pool MA, Weijers GWD, Ossenkoppele GJ and Feller N.
ISCT 2002; # 37.

Phenotypic expression of CD133 in acute leukemia

Expression of AC133 vs. CD34 in childhood acute leukemias. (article in German)

Ebener U, Brinkmann A, Zotova V, Niegemann E, Wehner S.
Klin Pädiatr. 2000 May-Jun; 212(3): 90-98.

General interest and research on CD133 expression

A novel five-transmembrane haematopoietic stem cell antigen: isolation, characterization, and molecular cloning.

Miraglia S, Godfrey W, Yin AH, Atkins K, Warnke R, Holden JT, Bray RA, Waller EK, Buck DW.
Blood. 1997 Dec 15; 90(12): 5013-5021.

AC133, a novel marker for human haematopoietic stem and progenitor cells.

Yin AH, Miraglia S, Zanjani ED, Almeida-Porada G, Ogawa M, Leary AG, Olweus J, Kearney J, Buck DW.
Blood. 1997 Dec 15; 90(12):5002-5012.

Clonogenicity and functionality of CD133 cells

Mesenchymal stem cells derived from CD133-positive cells in mobilized peripheral blood and cord blood: proliferation, Oct4 expression, and plasticity.

Tondreau T, Meuleman N, Delforge A, Dejeneffe M, Leroy R, Massy M, Mortier C, Bron D, Lagneaux L.
Stem Cells. 2005 Sep; 23(8):1105-12.

Biology and plasticity of CD133+ hematopoietic stem cells.

Handgretinger R, Gordon PR, Leimig T, Chen X, Buhring HJ, Niethammer D, Kuci S.
Ann N Y Acad Sci. 2003 May; 996:141-151.

Identification of a novel class of human adherent CD34- stem cells that give rise to SCID repopulating cells.

Kuci S, Wessels JT, Buhring HJ, Schilbach K, Schumm M, Seitz G, Loffler J, Bader P, Schlegel PG, Niethammer D, Handgretinger R.
Blood 2003 Feb 1; 101(3):869-876.

AC133+ umbilical cord blood progenitors demonstrate rapid self-renewal and low apoptosis.

Forraz N, Pettengell R, Deglesne PA, McGuckin CP.
Br J Haematol. 2002 Nov; 119(2):516-24.

Normal human bone marrow CD34+CD133+ cells contain primitive cells able to produce different categories of colony-forming unit megakaryocytes in vitro.

Charrier S, Boiret N, Fouassier M, Berger J, Rapatel C, Pigeon P, Mareynat G, Bonhomme J, Camilleri L, Berger MG.
Exp Hematol. 2002 Sep; 30(9):1051-60.

In vitro proliferative potential of AC133 positive cells in peripheral blood

Matsumoto K, Yasui K, Yamashita N, Horie Y, Yamada T, Tani Y, Shibata H, Nakano T, et al
Stem Cells. 2000; 18: 196-203.

A functional hierarchy among the CD34⁺ hematopoietic cells based on in vitro proliferative and differentiative potential of AC133⁺ CD34^{bright} and AC133^{dim}-CD34⁺ human cord blood cells.

Goussetis E, Theodosaki M, Paterakis G, Peristeri J, Petropoulos D, Kitra V, Papassarandis C, Graphakos S.

J Hematother Stem Cell Res. 2000 Dec; 9(6):827-840.

CD34⁺AC133⁺ cells isolated from cord blood are highly enriched in long-term culture-initiating cells, NOD/SCID-repopulating cells and dendritic cell progenitors.

de Wynter EA, Buck D, Hart C, Heywood R, Coutinho LH, Clayton A, Rafferty JA, Burt D, Guenechea G, Bueren JA, Gagen D, Fairbairn LJ, Lord BI, Testa NG.

Stem Cells. 1998; 16(6): 387-396.