

Reference list

PepTivator[®] Peptide Pools

Icheva, V. *et al.* (2013) Adoptive transfer of Epstein-Barr virus (EBV) nuclear antigen 1-specific T cells as treatment for EBV reactivation and lymphoproliferative disorders after allogeneic stem cell transplantation. *J. Clin. Oncol.* 31: 39–48.

Samuel, E.R. *et al.* (2013) Successful isolation and expansion of CMV-reactive T cells from G-CSF mobilized donors that retain a strong cytotoxic effector function. *Br. J. Haematol.* 160: 87–100.

Daenthanasamaka, A. *et al.* (2012) Integrase-defective lentiviral vectors encoding cytokines induce differentiation of human dendritic cells and stimulate multivalent immune responses *in vitro* and *in vivo*. *Vaccine* 30: 5118–5131.

Lugthart, G. *et al.* (2012) Simultaneous generation of multivirus-specific and regulatory T cells for adoptive immunotherapy. *J. Immunother.* 35: 42–53.

Guérin-El Khourouj, V. *et al.* (2011) Quantitative and qualitative CD4 T cell immune responses related to adenovirus DNAemia in hematopoietic stem cell transplantation. *Biol. Blood Marrow Transplant.* 17: 476–485.

Peggs, K.S. *et al.* (2011) Directly selected cytomegalovirus-reactive donor T cells confer rapid and safe systemic reconstitution of virus-specific immunity following stem cell transplantation. *Clin. Infect. Dis.* (2011) 52: 49–57.

Bes, M. *et al.* (2011) Reversal of non-structural protein 3-specific CD4⁺ T cell dysfunction in patients with persistent hepatitis C virus infection. *J. Viral Hepat.* 19: 283–294.

Beck, B. *et al.* (2011) Effects of TLR agonists on maturation and function of 3-day dendritic cells from AML patients in complete remission. *J. Transl. Med.* 9: 151.

Aïssi-Rothé, L. *et al.* (2010) Rapid generation of full clinical-grade human antiadenovirus cytotoxic T cells for adoptive immunotherapy. *J. Immunother.* 33: 414–424.

Draper, S. *et al.* (2010) Enhancing blood-stage malaria subunit vaccine immunogenicity in Rhesus macaques by combining adenovirus, poxvirus, and protein-in-adjuvant vaccines. *J. Immunol.* 185: 7583–7595.

Faist, B. *et al.* (2010) Cytomegalovirus infection- and age-dependent changes in human CD8 T cell cytokine expression patterns. *Clin. Vaccine Immunol.* 17: 986–992.

Lenders, K. *et al.* (2010) The effect of apoptotic cells on virus-specific immune responses detected using IFN-gamma ELISPOT. *J. Immunol. Methods* 357: 51–54.

Zandvliet, M. *et al.* (2010) Co-ordinated isolation of CD8 and CD4 T cells recognizing a broad repertoire of cytomegalovirus pp65 and IE1 epitopes for highly specific adoptive immunotherapy. *Cytotherapy* 12: 933–944.

Dodero A. *et al.* (2009) Haploidentical stem cell transplantation after a reduced intensity conditioning regimen for the treatment of advanced hematologic malignancies: posttransplantation CD8-depleted donor lymphocyte infusions contribute to improve T cell recovery. *Blood* 113: 4771–4779.

Miltenyi Biotec provides products and services worldwide. Visit www.miltenyibiotec.com/local to find your nearest Miltenyi Biotec contact.

MACS and PepTivator are registered trademarks of Miltenyi Biotec GmbH. Unless otherwise specifically indicated, Miltenyi Biotec products and services are for research use only and not for therapeutic or diagnostic use. Copyright © 2013 Miltenyi Biotec GmbH. All rights reserved.