

Contents

1. Description
 - 1.1 Background information
 - 1.2 Applications
2. Recommendations for *in vitro* restimulation of antigen-specific T cells with PepTivator® Peptide Pools
 - 2.1 Cell preparation
 - 2.2 Reagent requirements
 - 2.3 Recommendations for reconstitution of PepTivator® Peptide Pools
 - 2.4 Recommendations for *in vitro* restimulation of antigen-specific T cells
3. Appendix: Flask and dish sizes for *in vitro* T cell stimulation

1. Description

Components	6 nmol/peptide PepTivator® HCV1a NS5 – research grade, human: Pool of lyophilized peptides, consisting mainly of 15-mer sequences with 11 amino acids (aa) overlap, covering the sequence of the hepatitis C virus (HCV) genotype 1a NS5 protein (UniProtKB Acc. no. P26664 [aa1973-3011]).
Capacity	6 nmol (approximately 10 µg) per peptide for stimulation of up to 10 ⁸ total cells.
Product format	Lyophilized peptides containing stabilizer.
Purity	Average purity of peptides >70% (HPLC).
Storage	Store lyophilized product at –20 °C. The expiration date is indicated on the vial label.

This product contains no preservative and is sterile filtered; always handle under aseptic conditions.

1.1 Background information

Hepatitis C virus (HCV) is a single-stranded, enveloped RNA virus. After primary exposure to the virus most individuals develop a chronic viral hepatitis. HCV is the major cause of liver cirrhosis and hepatocellular carcinoma. The development of an immunological response to HCV is important for clearance of the acute infection and the long-term outcome of the pathogenesis in persistent infections.

The NS5 protein of HCV is a main target for CD4⁺ and CD8⁺ T cell responses.

PepTivator® Peptide Pools have been specially developed for efficient *in vitro* stimulation of antigen-specific CD4⁺ and CD8⁺ T cells, as peptides of 15 amino acid length with 11 amino acid overlap represent the optimized solution for stimulating both CD4⁺ and CD8⁺ T cells in various applications. Stimulation of

T cells with PepTivator Peptide Pools causes the secretion of effector cytokines and upregulation of activation markers, which then allows the detection or isolation of antigen-specific T cells. Quantitative, phenotypical, or functional analysis of antigen-specific T cell immunity can provide important information on the natural course of immune responses.

1.2 Applications

- Detection and analysis of antigen-specific CD4⁺ and CD8⁺ effector/memory T cells, for example, in PBMCs, by MACS® Cytokine Secretion Assays, intracellular cytokine staining, or other technologies.
- Isolation of viable antigen-specific CD4⁺ T cells with the CD154 MicroBead Kit.
- Isolation of viable antigen-specific CD4⁺ and CD8⁺ T cells using MACS Cytokine Secretion Assay – Cell Enrichment and Detection Kits or the CD137 MicroBead Kit for *in vitro* generation of T cell lines/clones.
- Generation of antigen-specific CD4⁺ and CD8⁺ effector/memory T cells from naive T cell populations for research on immunotherapy and vaccination.
- Pulsing of antigen-presenting cells for research on dendritic cell vaccination.

2. Recommendations for *in vitro* restimulation of antigen-specific T cells with PepTivator® Peptide Pools

2.1 Cell preparation

For induction of cytokine secretion by antigen-specific T cells, best results are achieved by stimulation of fresh PBMCs, whole blood, or other leukocyte-containing single-cell preparations from tissues or cell lines. Alternatively, frozen cell preparations can be used.

▲ **Note:** Remove platelets after density gradient separation. Resuspend cell pellet, fill tube with buffer, and mix. Centrifuge at 200×g for 10–15 minutes at 20 °C. Carefully remove supernatant.

▲ **Note:** PBMCs may be stored overnight. The cells should be resuspended and incubated in culture medium as described in 2.4, steps 1–3, but without addition of antigen. The antigen is then added to the culture on the next day.

For details about cell preparation refer to the protocols section at www.miltenyibiotec.com/protocols.

2.2 Reagent requirements

- Culture medium, e.g., RPMI 1640 (# 130-091-440) containing 5% human serum, e.g., autologous or AB serum.
- ▲ **Note:** Do not use bovine serum albumin (BSA) or fetal bovine serum (FBS) because of non-specific stimulation.

- (Optional) Cytokine Secretion Assay Kit. For additional reagent and instrument requirements refer to the data sheet of the respective Cytokine Secretion Assay.
- (Optional) Intracellular cytokine staining, e.g., with Anti-IFN- γ -PE (# 130-091-653). For additional reagent requirements refer to the respective data sheet. For more information on other fluorochrome-conjugates refer to www.miltenyibiotec.com.
- (Optional) Intracellular cytokine staining of activated CD4⁺ T cells by using, for example, the CD154/IFN- γ /CD4 Detection Kit (# 130-092-814).
- (Optional) CD154 MicroBead Kit (# 130-092-658). For details refer to the CD154 MicroBead Kit data sheet.
- (Optional) CD137 MicroBead Kit (# 130-093-476). For details refer to the CD137 MicroBead Kit data sheet.
- (Optional) CytoStim for restimulation of human T cells (# 130-092-172, # 130-092-173). For details refer to the CytoStim data sheet.

2.3 Recommendations for reconstitution of PepTivator® Peptide Pools

- For reconstitution of the lyophilized peptide pool take the vial from -20°C and warm-up to room temperature.
▲ **Note:** Do not open the vial by removing the rubber plug.
- To dissolve the 6 nmol PepTivator® Peptide Pool fill a sterile syringe (0.5 mL) with 200 μL of sterile water.
- Slowly inject the water with a sterile needle through the center of the rubber plug into the vial containing the lyophilized peptide pool.
- Vortex the solution to completely dissolve the lyophilized peptide pool.
The concentration of the stock solution of PepTivator Peptides is 30 nmol (approximately 50 μg) of each peptide per mL.
- Remove the rubber plug and aspirate the stock solution with a pipette.
- To avoid repeated freeze-thaw cycles prepare working aliquots from the stock solution.
- Store the working aliquots at -80°C .

2.4 Recommendations for *in vitro* restimulation of antigen-specific T cells

▲ HCV-specific T cells are expected to be present only in certain individuals. Their frequency may be low compared to T cells with other specificities. The given protocol for *in vitro* T cell stimulation thus may only serve as a guideline.

▲ Magnetic enrichment of stimulated antigen-specific T cells according to cytokine secretion using the MACS Secretion Assay Technology or according to expression of activation marker, e.g. CD154, will enhance the sensitivity to detect rare cells.

▲ Always include a negative control (without antigen) in the experiment. As a positive control, a sample stimulated with, e.g. CytoStim, may also be included.

- Wash cells by adding medium, centrifuge at 300 \times g for 10 minutes. Aspirate supernatant.
- Resuspend cells in culture medium at 10^7 cells/mL. Plate cells

in dishes at a density of 5×10^6 cells/cm² (refer to 3. Appendix: Flask and dish sizes for *in vitro* T cell stimulation).

- Mix the reconstituted PepTivator thoroughly. Add 20 μL of PepTivator stock solution per mL cell suspension. Mix carefully and incubate cells at 37°C and 5% CO₂.

The final concentration of PepTivator in the cell suspension is 0.6 nmol (approximately 1 μg) of each peptide/mL.

Cytokine Secretion Assay: Incubate cells for 3–6 hours.

CD154 MicroBead Kit: Incubate cells for 4–16 hours.

CD137 MicroBead Kit: Incubate cells for 16–24 hours.

Intracellular cytokine staining antibodies or kits, e.g., CD154/IFN- γ /CD4 Detection Kit: Incubate cells for 2 hours, then add 1 $\mu\text{g}/\text{mL}$ brefeldin A, and incubate for further 4 hours.

- Collect cells carefully by using a cell scraper, or by pipetting up and down when working with smaller volumes. Rinse the dish with cold buffer. Check microscopically for any remaining cells, if necessary, rinse the dish again.

To proceed with the Cytokine Secretion Assay, the CD154 or CD137 MicroBead Kits, or intracellular cytokine staining, please refer to the respective data sheet.

▲ **Note:** When preparing cells for **intracellular cytokine staining**, fixed cells may be stored at $2-8^{\circ}\text{C}$ for up to 1 week.

3. Appendix: Flask and dish sizes for *in vitro* T cell stimulation

For *in vitro* T cell stimulation (refer to 2.4) the cells should be resuspended in culture medium, containing 5% of human serum, at a dilution of 10^7 cells/mL. The cells should be plated at a density of 5×10^6 cells/cm². Both the dilution and the cell density are important to assure optimum stimulation.

The following table lists culture plate, dish and flask sizes suitable for different cell numbers. It also indicates the appropriate amount of medium to add.

Total cell number	Medium volume to add	Culture plate	Well diameter
0.15×10^7	0.15 mL	96 well	0.64 cm
0.50×10^7	0.50 mL	48 well	1.13 cm
1.00×10^7	1.00 mL	24 well	1.60 cm
2.00×10^7	2.00 mL	12 well	2.26 cm
5.00×10^7	5.00 mL	6 well	3.50 cm
Total cell number	Medium volume to add	Culture dish	Dish diameter
4.5×10^7	4.5 mL	small	3.5 cm
10.0×10^7	10.0 mL	medium	6 cm
25.0×10^7	25.0 mL	large	10 cm
50.0×10^7	50.0 mL	extra large	15 cm
Total cell number	Medium volume to add	Culture flask	Growth area
12×10^7	12 mL	50 mL	25 cm ²
40×10^7	40 mL	250 mL	75 cm ²
80×10^7	80 mL	720 mL	162 cm ²
120×10^7	120 mL	900 mL	225 cm ²

Refer to www.miltenyibiotec.com for all data sheets and protocols. Miltenyi Biotec provides technical support worldwide. Visit www.miltenyibiotec.com/local to find your nearest Miltenyi Biotec contact.

Legal notices

Limited product warranty

Miltenyi Biotec B.V. & Co. KG and/or its affiliate(s) warrant this product to be free from material defects in workmanship and materials and to conform substantially with Miltenyi Biotec's published specifications for the product at the time of order, under normal use and conditions in accordance with its applicable documentation, for a period beginning on the date of delivery of the product by Miltenyi Biotec or its authorized distributor and ending on the expiration date of the product's applicable shelf life stated on the product label, packaging or documentation (as applicable) or, in the absence thereof, ONE (1) YEAR from date of delivery ("Product Warranty"). Miltenyi Biotec's Product Warranty is provided subject to the warranty terms as set forth in Miltenyi Biotec's General Terms and Conditions for the Sale of Products and Services available on Miltenyi Biotec's website at www.miltenyibiotec.com, as in effect at the time of order ("Product Warranty"). Additional terms may apply. BY USE OF THIS PRODUCT, THE CUSTOMER AGREES TO BE BOUND BY THESE TERMS.

THE CUSTOMER IS SOLELY RESPONSIBLE FOR DETERMINING IF A PRODUCT IS SUITABLE FOR CUSTOMER'S PARTICULAR PURPOSE AND APPLICATION METHODS.

Technical information

The technical information, data, protocols, and other statements provided by Miltenyi Biotec in this document are based on information, tests, or experience which Miltenyi Biotec believes to be reliable, but the accuracy or completeness of such information is not guaranteed. Such technical information and data are intended for persons with knowledge and technical skills sufficient to assess and apply their own informed judgment to the information. Miltenyi Biotec shall not be liable for any technical or editorial errors or omissions contained herein.

All information and specifications are subject to change without prior notice. Please contact Miltenyi Biotec Technical Support or visit www.miltenyibiotec.com for the most up-to-date information on Miltenyi Biotec products.

Licenses

This product and/or its use may be covered by one or more pending or issued patents and/or may have certain limitations. Certain uses may be excluded by separate terms and conditions. Please contact your local Miltenyi Biotec representative or visit Miltenyi Biotec's website at www.miltenyibiotec.com for more information.

The purchase of this product conveys to the customer the non-transferable right to use the purchased amount of the product in research conducted by the customer (whether the customer is an academic or for-profit entity). This product may not be further sold. Additional terms and conditions (including the terms of a Limited Use Label License) may apply.

CUSTOMER'S USE OF THIS PRODUCT MAY REQUIRE ADDITIONAL LICENSES DEPENDING ON THE SPECIFIC APPLICATION. THE CUSTOMER IS SOLELY RESPONSIBLE FOR DETERMINING FOR ITSELF WHETHER IT HAS ALL APPROPRIATE LICENSES IN PLACE. Miltenyi Biotec provides no warranty that customer's use of this product does not and will not infringe intellectual property rights owned by a third party. BY USE OF THIS PRODUCT, THE CUSTOMER AGREES TO BE BOUND BY THESE TERMS.

Trademarks

MACS, the Miltenyi Biotec logo, and PepTivator are registered trademarks or trademarks of Miltenyi Biotec and/or its affiliates in various countries worldwide.

Copyright © 2020 Miltenyi Biotec and/or its affiliates. All rights reserved.