



Selected references

gentleMACS™ Dissociators and kits

Human tumor

Isolation of tumor-associated macrophages (TAM)

Hoppstädter, J. *et al.* (2015) M2 polarization enhances silica nanoparticle uptake by macrophages. *Front. Pharmacol.* 6: 55.

www.ncbi.nlm.nih.gov/pubmed/25852557

Isolation and expansion of TILs from renal cell carcinoma (RCC)

Baldan, V. *et al.* (2015) Efficient and reproducible generation of tumour-infiltrating lymphocytes for renal cell carcinoma. *Br. J. Cancer* 112: 1510–1518.

www.ncbi.nlm.nih.gov/pubmed/25867267

Isolation of tumor cells from bladder tumors

Naipal, K. *et al.* (2015) Attenuated XPC expression is not associated with impaired DNA repair in bladder cancer. *PLoS One* 10: e0126029.

www.ncbi.nlm.nih.gov/pubmed/25927440

Isolation and analysis of tumor cells and pancreatic stellate cells from pancreatic tumor

Fujiwara, K. *et al.* (2014) CD166/ALCAM expression is characteristic of tumorigenicity and invasive and migratory activities of pancreatic cancer cells. *PLoS One* 9: e107247.

www.ncbi.nlm.nih.gov/pubmed/25221999

Isolation and analysis of tumor-infiltrating Tregs from colorectal and lung tumor (NSCLC)

De Simone, M. *et al.* (2016) Transcriptional landscape of human tissue lymphocytes unveils uniqueness of tumor-infiltrating T regulatory cells. *Immunity* 45: 1135–1147.

www.ncbi.nlm.nih.gov/pubmed/27851914

Analysis of tumor-associated macrophages (TAMs) from glioblastoma

Castro, B.A. *et al.* (2017) Macrophage migration inhibitory factor downregulation: a novel mechanism of resistance to anti-angiogenic therapy. *Oncogene*. doi: 10.1038/onc.2017.

www.ncbi.nlm.nih.gov/pubmed/28218903

Xenograft tumor

Isolation of tumor cells and depletion of mouse cells from breast cancer patient-derived xenografts

Aloia, A. *et al.* (2015) The sialyl-glycolipid stage-specific embryonic antigen 4 marks a subpopulation of chemotherapy-resistant breast cancer cells with mesenchymal features. *Breast Cancer Res.* 17: 146.

www.ncbi.nlm.nih.gov/pubmed/26607327

Isolation and culture of primary tumor cells from human gliomas

Parajuli, P. *et al.* (2015) Preferential expression of functional IL-17R in glioma stem cells: potential role in self-renewal. *Oncotarget* 7: 6121–6135.

www.ncbi.nlm.nih.gov/pubmed/26755664

Depletion of mouse cells from human tumor xenografts for next generation sequencing (NGS)

Agorku, D.J. *et al.* (2016) Depletion of mouse cells from human tumor xenografts significantly improves downstream analysis of target cells. *J. Vis. Exp.* 113: doi:10.3791/54259.

www.ncbi.nlm.nih.gov/pubmed/27501218

Isolation of tumor cells from colorectal cancer patient-derived xenografts

Arena, S. *et al.* (2016) MM-151 overcomes acquired resistance to cetuximab and panitumumab in colorectal cancers harboring EGFR extracellular domain mutations. *Sci. Transl. Med.* 8: 324ra14.

www.ncbi.nlm.nih.gov/pubmed/26843189

High-throughput isolation of tumor cells and depletion of mouse cells from breast cancer patient-derived xenografts

Chen, Y.-C. *et al.* (2016) High-throughput single-cell derived sphere formation for cancer stem-like cell identification and analysis. *Sci. Rep.* 6: 27301.

www.ncbi.nlm.nih.gov/pubmed/27292795

Isolation and analysis of tumor-infiltrating MDSCs and T cells from mouse breast cancer model and patient-derived xenografts

Welte, T. *et al.* (2016) Oncogenic mTOR signalling recruits myeloid-derived suppressor cells to promote tumour initiation. *Nat. Cell Biol.* 18: 632–644.

www.ncbi.nlm.nih.gov/pubmed/27183469

Isolation and genomic analysis of tumor cells after depletion of mouse cells from small-cell lung cancer patient-derived xenografts

Schneeberger, V.E. *et al.* (2016) Quantitation of murine stroma and selective purification of the human tumor component of patient-derived xenografts for genomic analysis. *PLoS One* 11: e0160587.

www.ncbi.nlm.nih.gov/pubmed/27611664

Isolation and analysis of cancer stem cells from colorectal cancer xenografts

Oshima, N. *et al.* (2014) Induction of cancer stem cell properties in colon cancer cells by defined factors. *PLoS One* 9: e101735.

www.ncbi.nlm.nih.gov/pubmed/25006808

Mouse tumor

Flow and transcriptome analysis of tumor-infiltrating myeloid-derived cells from synovial sarcoma

Barrott, J. *et al.* (2016) Modeling synovial sarcoma metastasis in the mouse: PI3'-lipid signaling and inflammation. *J. Exp. Med.* 213: 2989–3005.

www.ncbi.nlm.nih.gov/pubmed/27956588

Flow analysis of tumor-infiltrating lymphocytes from B16F10 melanoma

Chen, S. *et al.* (2014) Combination of 4-1BB agonist and PD-1 antagonist promotes antitumor effector/memory CD8 T cells in a poorly immunogenic tumor model. *Cancer Immunol. Res.* 3: 149–160.

www.ncbi.nlm.nih.gov/pubmed/25387892

Flow analysis and RNA isolation of tumor-infiltrating MDSCs and TILs from TC-1 tumors

Mikyšková, R. *et al.* (2014) DNA demethylating agent 5-azacytidine inhibits myeloid-derived suppressor cells induced by tumor growth and cyclophosphamide treatment. *J. Leukoc. Biol.* 95: 1–11.

www.ncbi.nlm.nih.gov/pubmed/24389335

Flow analysis of tumor-infiltrating MDSCs and TILs from thymoma

Qin, H. *et al.* (2014) Generation of a new therapeutic peptide that depletes myeloid-derived suppressor cells in tumor-bearing mice. *Nat. Med.* 20: 676–681.

www.ncbi.nlm.nih.gov/pubmed/24859530

Isolation of cancer stem cells from glioblastoma

Pessina, S. *et al.* (2016) The multidrug-resistance transporter Abcc3 protects NK cells from chemotherapy in a murine model of malignant glioma. *Oncoimmunology* 5: e1108513.

www.ncbi.nlm.nih.gov/pubmed/27467914

Lamina propria

Isolation of dendritic cells from mouse colon

Nakahashi-Oda, C. *et al.* (2016) Apoptotic epithelial cells control the abundance of Treg cells at barrier surfaces. *Nat. Immunol.* 17: 441–450.

www.ncbi.nlm.nih.gov/pubmed/26855029

Isolation of innate lymphoid cells from mouse small intestine

Paclik, D. *et al.* (2015) ICOS regulates the pool of group 2 innate lymphoid cells under homeostatic and inflammatory conditions in mice. *Eur. J. Immunol.* 45: 2766–2772.

www.ncbi.nlm.nih.gov/pubmed/26249010

Isolation of intestinal cells from pups' small intestine

Schulz, S. *et al.* (2013) Heme oxygenase-1 deficiency promotes the development of necrotizing enterocolitis-like intestinal injury in a newborn mouse model. *Am. J. Physiol. Gastrointest. Liver Physiol.* 304: G991–G1001.

www.ncbi.nlm.nih.gov/pubmed/23578787

Lung

RNA extraction from small intestine and lung

Platzer, B. *et al.* (2015) Dendritic cell-bound IgE functions to restrain allergic inflammation at mucosal sites. *Mucosal Immunol.* 8: 516–532.

www.ncbi.nlm.nih.gov/pubmed/25227985

Isolation of IL-7-producing cells from mouse lung

Shinoda, K. *et al.* (2016) Thy1⁺IL-7⁺ lymphatic endothelial cells in iBALT provide a survival niche for memory T-helper cells in allergic airway inflammation. *Proc. Natl. Acad. Sci USA* 113: E2842–2851.

www.ncbi.nlm.nih.gov/pubmed/27140620

Isolation of mast cell progenitors from mouse lung

Dahlin, J.S. *et al.* (2015) Distinguishing mast cell progenitors from mature mast cells in mice. *Stem Cells Dev.* 24: 1703–1711.

www.ncbi.nlm.nih.gov/pubmed/25744159

Isolation of influenza virus from frozen mouse lung

Moorthy, A.N. *et al.* (2016) Effect of high-fat diet on the formation of pulmonary neutrophil extracellular traps during influenza pneumonia in BALB/c mice. *Front. Immunol.* 7: 289.2

www.ncbi.nlm.nih.gov/pubmed/27531997

Adult brain

Isolation of living neonatal and adult astrocytes

Kantzer, C.G. *et al.* (2017) Anti-ACSA-2 defines a novel monoclonal antibody for prospective isolation of living neonatal and adult astrocytes. *Glia* 65: 990–1004.

www.ncbi.nlm.nih.gov/pubmed/28317180

Isolation of neurons from adult mouse brain

Berl, S. *et al.* (2017) Enrichment and isolation of neurons from adult mouse brain for *ex vivo* analysis. *J. Neurosci. Methods* 283: 15–22.

www.ncbi.nlm.nih.gov/pubmed/28336359

Isolation of cells from Alzheimer's disease (AD) model mice for extracellular vesicle analysis

Dinkins, M.B. *et al.* (2016) Neutral sphingomyelinase-2 deficiency ameliorates Alzheimer's disease pathology and improves cognition in the 5XFAD mouse. *J. Neurosci.* 36: 8653–8667.

www.ncbi.nlm.nih.gov/pubmed/27535912

Isolation of CD3⁺ T cells from adult mouse brain

Jarry, U. *et al.* (2016) Stereotaxic administrations of allogeneic human V γ 9V δ 2 T cells efficiently control the development of human glioblastoma brain tumors. *Oncoimmunology* 5: e1168554.

www.ncbi.nlm.nih.gov/pubmed/27471644

Neural tissue

Isolation of astrocytes and microglia from P30–P50 mouse brain

Cormier, S.A. *et al.* (2014) Limited type I interferons and plasmacytoid dendritic cells during neonatal respiratory syncytial virus infection permit immunopathogenesis upon reinfection. *J. Virol.* 88: 9350–9360.

www.ncbi.nlm.nih.gov/pubmed/24920801

Preparation of brain cortex tissue and isolation of endothelial cells from P1–P2 and adult mouse

Wylot, B. *et al.* (2015) Isolation of vascular endothelial cells from intact and injured murine brain cortex—technical issues and pitfalls in FACS analysis of the nervous tissue. *Cytometry A* 87: 908–920.

www.ncbi.nlm.nih.gov/pubmed/25892199

Isolation of astrocytes from subventricular zone of mouse brain

Llorens-Bobadilla, E. *et al.* (2015) Single-cell transcriptomics reveals a population of dormant neural stem cells that become activated upon brain injury. *Cell Stem Cell* 17: 329–340.

www.ncbi.nlm.nih.gov/pubmed/26235341

Isolation of astrocytes and neurons from P9–10 mouse brain

Feldmann, M. *et al.* (2014) Isolating astrocytes and neurons sequentially from postnatal murine brains with a magnetic cell separation technique. *J. Biol. Methods* 1: e11.

www.jbmethods.org/jbm/article/view/33

Isolation of hippocampal cells from hippocampus tissue

Cohen, S. *et al.* (2016) Adverse early life environment increases hippocampal microglia abundance in conjunction with decreased neural stem cells in juvenile mice. *Int. J. Dev. Neurosci.* 55: 56–65.

www.ncbi.nlm.nih.gov/pubmed/27666383

Liver

Isolation of macrophages and NK cells from mouse liver

Kouwaki, T. *et al.* (2016) Extracellular vesicles including exosomes regulate innate immune responses to hepatitis B virus infection. *Front. Immunol.* 7: 335.

www.ncbi.nlm.nih.gov/pubmed/27630638

Protein extraction from frozen human liver

Descamps, V. *et al.* (2012) Strong correlation between liver and serum levels of hepatitis C virus core antigen and RNA in chronically infected patients. *J. Clin. Microbiol.* 50: 465–468.

www.ncbi.nlm.nih.gov/pubmed/22162563

Spleen

Isolation of pDCs from mouse spleen

Cormier, S.A. *et al.* (2014) Limited type I interferons and plasmacytoid dendritic cells during neonatal respiratory syncytial virus infection permit immunopathogenesis upon reinfection. *J. Virol.* 88: 9350–9360.

www.ncbi.nlm.nih.gov/pubmed/24920801

Homogenization of human spleen

Martín-Jaular, L. *et al.* (2016) [published erratum in *Front. Cell Dev. Biol.* 2017; 4: 153] Spleen-dependent immune protection elicited by CpG adjuvanted reticulocyte-derived exosomes from malaria infection is associated with changes in T cell subsets' distribution. *Front. Cell Dev. Biol.* 4: 1–11.

www.ncbi.nlm.nih.gov/pubmed/27900319

Neonatal heart

Isolation of ventricular myocytes from neonatal rat heart

Trantidou, T. *et al.* (2015) Biorealistic cardiac cell culture platforms with integrated monitoring of extracellular action potentials. *Sci. Rep.* 5: 11067.

www.ncbi.nlm.nih.gov/pubmed/26053434

Isolation and gene profiling of cardiomyocytes from embryonic and neonatal mouse heart

Wiencierz, A.M. *et al.* (2015) Differential expression levels of integrin $\alpha 6$ enable the selective identification and isolation of atrial and ventricular cardiomyocytes. *PLoS One* 10: e0143538.

www.ncbi.nlm.nih.gov/pubmed/26618511

Isolation of cardiomyocytes from neonatal mouse heart

Raulf, A. *et al.* (2015) Transgenic systems for unequivocal identification of cardiac myocyte nuclei and analysis of cardiomyocyte cell cycle status. *Basic Res. Cardiol.* 110: 33.

www.ncbi.nlm.nih.gov/pubmed/25925989

Skeletal muscle

Isolation of neutrophils from *C. perfringens*-infected femoral muscle

Takehara, M. *et al.* (2016) *Clostridium perfringens* α -toxin impairs innate immunity via inhibition of neutrophil differentiation. *Sci. Rep.* 6: 28192.

www.ncbi.nlm.nih.gov/pubmed/27306065

Isolation of satellite cells from mouse skeletal muscle

Boettger, T. *et al.* (2014) The miR-206/133b cluster is dispensable for development, survival and regeneration of skeletal muscle. *Skelet. Muscle* 4: 23.

www.ncbi.nlm.nih.gov/pubmed/25530839

Isolation and cultivation of satellite cells from mouse skeletal muscle

Musarò, A. and Carosio, S. (2017) Isolation and culture of satellite cells from mouse skeletal muscle. *Methods Mol. Biol.* 1553: 155–167.

www.ncbi.nlm.nih.gov/pubmed/28229414

Skin

Isolation of Langerhans cells from mouse epidermis

Nakahashi-Oda, C. *et al.* (2016) Apoptotic epithelial cells control the abundance of Treg cells at barrier surfaces. *Nat. Immunol.* 17: 441–450.

www.ncbi.nlm.nih.gov/pubmed/26855029

Cell isolation from Zika virus (ZIKV)-infected human skin biopsy

Hamel, R. *et al.* (2015) Biology of Zika virus infection in human skin cells. *J Virol.* 89: 8880–8896.

www.ncbi.nlm.nih.gov/pubmed/26085174

Isolation of skin-infiltrating cells from mouse whole skin

Okamoto, S. *et al.* (2015) Anti-IL-12/23 p40 antibody attenuates experimental chronic graft-versus-host disease via suppression of IFN- γ /IL-17-producing cells. *J. Immunol.* 194: 1357–1363.

www.ncbi.nlm.nih.gov/pubmed/25527789

Adipose tissue

Isolation of progenitor cells from stromal-vascular fraction (SVF) of mouse adipose tissue

Bayindir, I. *et al.* (2015) Transcriptional pathways in cPGI2-induced adipocyte progenitor activation for browning. *Front. Endocrinol. (Lausanne)* 6: 129.

www.ncbi.nlm.nih.gov/pubmed/26347713

Others

Homogenization of human lymph nodes

Tauriainen, J. *et al.* (2017) Perturbed CD8⁺ T cell TIGIT/CD226/PVR axis despite early initiation of antiretroviral treatment in HIV infected individuals. *Sci. Rep.* 7: 40354.

www.ncbi.nlm.nih.gov/pubmed/28084312

Homogenization of fecal pellets

Cheng, Y.C. *et al.* (2016) Early and non-invasive detection of chronic wasting disease prions in elk feces by real-time quaking induced conversion. *PLoS One* 11: 1–18.

www.ncbi.nlm.nih.gov/pubmed/27829062

Isolation of mitochondria

Cowan, D.B. *et al.* (2016) Intracoronary delivery of mitochondria to the ischemic heart for cardioprotection. *PLoS One* 11: 1–19.

www.ncbi.nlm.nih.gov/pubmed/27500955



Miltenyi Biotec provides products and services worldwide. Visit www.miltenyibiotec.com/local to find your nearest Miltenyi Biotec contact.

Unless otherwise specifically indicated, Miltenyi Biotec products and services are for research use only and not for therapeutic or diagnostic use. MACS, the MACS logo, and gentleMACS are registered trademarks or trademarks of Miltenyi Biotec GmbH and/or its affiliates in various countries worldwide. Copyright © 2017 Miltenyi Biotec GmbH and/or its affiliates. All rights reserved.